

The A -Module Structure Induced by a Drinfeld A -Module of Rank 2 over a Finite Field

MOHAMED AHMED Mohamed Saadbouh

Institut de Mathématiques de Luminy, CNRS-UPR 9016

Case 907, 163 Avenue de Luminy, 13288 Marseille, Cedex 9. France.

Résumé

Soit Φ un $\mathbf{F}_q[T]$ -module de Drinfeld de rang 2, sur un corps fini L , une extension de degré n d'un corps fini à q éléments $\mathbf{F}_q$. Soit $P_\Phi(X) = X^2 - cX + \mu P^m$ (c un élément de $\mathbf{F}_q[T]$, μ un élément non nul de $\mathbf{F}_q$, m le degré de l'extension L sur le corps $\mathbf{F}_q[T]/P$ et P est la $\mathbf{F}_q[T]$ -caractéristique de L) le polynôme caractéristique, de Frobenius F de L . On s'intéressera à la structure de $\mathbf{F}_q[T]$ -module fini L^Φ et on prouvera notre résultat principal qui est l'analogie du théorème de Deuring pour les courbes elliptiques : soit $M = \frac{\mathbf{F}_q[T]}{I_1} \oplus \frac{\mathbf{F}_q[T]}{I_2}$, où $I_1 = (i_1)$, $I_2 = (i_2)$ (i_1, i_2 deux polynômes de $\mathbf{F}_q[T]$) et tel que : $i_2 \mid (c-2)$. Alors il existe un $\mathbf{F}_q[T]$ -module de Drinfeld Φ sur L de rang 2, ordinaire, tel que : $L^\Phi \simeq M$. On fini par une statistique concernant la cyclicité de la structure de A -module L^Φ .

Abstract

Let Φ be a Drinfeld $\mathbf{F}_q[T]$ -module of rank 2, over a finite field L , a finite extension of n degrees for a finite field with q elements $\mathbf{F}_q$. Let $P_\Phi(X) = X^2 - cX + \mu P^m$ (c an element of $\mathbf{F}_q[T]$ and μ a no null element of $\mathbf{F}_q$, m the degree of the extension L over the field $\mathbf{F}_q[T]/P$, P is a $\mathbf{F}_q[T]$ -characteristic of L and d the degree of the polynomial P) the characteristic polynomial, of the Frobenius F of L . We will interested to the structure of finite $\mathbf{F}_q[T]$ -module L^Φ deduct by Φ over L and will proof our main result, the analogue of Deuring theorem for the elliptic curves : Let $M = \frac{\mathbf{F}_q[T]}{I_1} \oplus \frac{\mathbf{F}_q[T]}{I_2}$, where $I_1 = (i_1)$, $I_2 = (i_2)$ (i_1, i_2 two polynomials of $\mathbf{F}_q[T]$) and such that : $i_2 \mid (c-2)$. Then there exists an ordinary Drinfeld $\mathbf{F}_q[T]$ -module Φ over L of rank 2, such that : $L^\Phi \simeq M$. We finish by a statistic about the cyclicity of such structure L^Φ , and we prove that is cyclic only for the trivial extensions of $\mathbf{F}_q$.

1 Introduction

Let E be an elliptic curve over finite field $\mathbf{F}_q$, we know, see [12], [15], [16] and [19], that the endomorphism ring of E , $\text{End}_{\mathbf{F}_q} E$, is an $\mathbb{Z}$ -order in a division

algebra, this algebra is : $\mathbb{Q}$ and in this case $\text{End}_{\mathbf{F}_q} E = \mathbb{Z}$, or a quadratic complex field and in this case : $\text{End}_{\mathbf{F}_q} E = \mathbb{Z} + c O_K$ where c is an element of $\mathbb{Z}$ and O_K is the maximal $\mathbb{Z}$ -order in this quadratic complex field, or is a Quaternion Algebra over $\mathbb{Q}$ and in this case $\text{End}_{\mathbf{F}_q} E$ is a maximal order in this Quaternion Algebra. We put $E(\mathbf{F}_q)$ the abelian group of $\mathbf{F}_q$ -rational points of E . The cardinal of this abelian group is equal to $N = q + 1 - c$, and by Hasse-Weil $|c| \leq 2\sqrt{q}$. The structure of this group in the ordinary case is :

$$E(\mathbf{F}_q) \simeq \mathbb{Z}/A \oplus \mathbb{Z}/B, \text{ if } (c, q) = 1, \quad B \mid A, \quad B \mid (c - 2) \quad \text{and} \quad A.B = N \quad (1)$$

conversely, for every abelian group $\mathbb{Z}/A \oplus \mathbb{Z}/B$, with $B \mid A$, $B \mid (c - 2)$ and $A.B = N$, there is an elliptic curve E such that : $E(\mathbf{F}_q) \simeq \mathbb{Z}/A \oplus \mathbb{Z}/B$, we note that in the supersingular case, this structure is also known, see [15]. Over this structure, S. Vladut, in [20], has effected a statistic about the report of elliptic curves for which the $E(\mathbf{F}_q)$ is cyclic over the number of classes of $\mathbf{F}_q$ -endomorphisms of elliptic curves over finite field $\mathbf{F}_q$, this report will depend on q and will be noted $c(q)$, and we have :

$$c(q) = \frac{\#\{E, E(\mathbf{F}_q) \text{ cyclic}\}}{\#\{E\}},$$

where $\#\{E\}$ is the number of classes of $\mathbf{F}_q$ -endomorphisms of elliptic curve over a finite field $\mathbf{F}_q$, and we know, see [20], that :

$c(q) = 1$ if and only if $q = 2^l$ where $l \neq 2$ is a prime or equal at 1 and one of the following conditions is satisfied:

1. $q - 1$ is prime, $q \neq 4$ (the case $q = 2$ is included, thus we consider 1 as prime),
2. $q - 1 = l_1 l_2$ with primes l_1 and l_2 is not a " small " divisors of $q - 1$;
3. $q - 1 = l_1 l_2 l_3$ with primes l_1, l_2 and l_3 are not a " small " divisors of $q - 1$.

the case $l_1 = l_2$ is not exclude.

In general case, the number $c(q)$ is given in [20], by :

Let $\varepsilon > 0$ we have:

$$c(q) = \prod_l \left(1 - \frac{1}{l(l^2 - 1)}\right) + O(q^{-1/2+\varepsilon}),$$

where the product is taken over all prime divisors of $q - 1$. Our goal here is to give an analog of the above mentioned results in the case of Drinfeld Modules of rank 2. We recall quickly what is it : let K a no empty global field of characteristic p (that means a rational functions field of one indeterminate over a finite field) with a constant field the finite field $\mathbf{F}_q$ with p^s elements. We fix one place of K , noted ∞ and we call A the ring of regular elements away

from the place ∞ . Let L be a commutative field of characteristic p , and let $\gamma : A \rightarrow L$, be a A -ring homomorphism, the kernel of this homomorphism is noted P and $m = [L, A/P]$ is the extension degrees of L over A/P .

We note $L\{\tau\}$ the Ore's polynomial ring, that means the polynomial ring of τ , τ is the Frobenius of $\mathbf{F}_q$, with the usual addition and the product is given by the computation rule : for every λ of L , $\tau\lambda = \lambda^q\tau$. We say a Drinfeld A -module Φ for a non trivial homomorphism of ring, from A to $L\{\tau\}$ which is different of γ . this homomorphism Φ , once defined, gives a A -module structure over the A -field L , noted L^Φ , where the name of a Drinfeld A -module for a homomorphism Φ . This structure A -module is depending on Φ and especially on this rank.

Let χ be the characteristic of Euler-Poincare (it is a ideal from A), so we can speak about the ideal $\chi(L^\Phi)$, will be noted by χ_Φ , which is by Definition a divisor for A , corresponding for the elliptic curves to a number of points of the variety over their basic fields.

We will work, in this paper, in the special case $K = \mathbf{F}_q(T)$, $A = \mathbf{F}_q[T]$. Let $P_\Phi(X)$: the characteristic polynomial of the A -module Φ , it is also a characteristic polynomial of Frobenius F of L . We can prove that this polynomial can be given by : $P_\Phi(X) = X^2 - cX + \mu P^m$, such that $\mu \in \mathbf{F}_q^*$, $c \in A$ and $\deg c \leq m \cdot \frac{d}{2}$, the Hass-Weil's analogue in this case.

We will interest to a Drinfeld A -module structure L^Φ in the case of rank 2, and we will prove that for an ordinary Drinfeld $\mathbf{F}_q[T]$ -module, this structure is always the sum of two cyclic and finite $\mathbf{F}_q[T]$ -modules : $\frac{A}{I_1} \oplus \frac{A}{I_2}$ where $I_1 = (i_1)$ and $I_2 = (i_2)$ such that i_1 and i_2 is two ideals of A , which verifies $i_2 \mid i_1$. We will show that $\chi_\Phi = I_1 I_2 = (P_\Phi(1))$, and if we put $i = \text{paced}(i_1, i_2)$, then : $i^2 \mid P_\Phi(1)$. We will give now some appears of our results proved in this paper :

Proposition 1.1. *With the above notations, we have :*

$L^\Phi \simeq \frac{A}{I_1} \oplus \frac{A}{I_2}$. And if we have an ordinary module Φ , then : $i_2 \mid (c - 2)$.

We note by $\text{End}_L \Phi$ the endomorphism ring of a Drinfeld A -module Φ , we have:

Proposition 1.2. *Let Φ be an ordinary Drinfeld A -module of rank 2 and let ρ a prime ideal of A different from P the A -characteristic of L , such that $\rho^2 \mid P_\Phi(1)$ and $\rho \mid (c - 2)$. Then $\Phi(\rho) \subset L^\Phi$ if and only if the A -order $O(\Delta/\rho^2) \subset \text{End}_L \Phi$.*

Finally, we come to our main result, which is a complete analog of (1), the Deuring-Waterhous theorem for the elliptic curves :

Theorem 1.1. *Let $M = \frac{A}{I_1} \oplus \frac{A}{I_2}$, where $I_1 = (i_1)$, $I_2 = (i_2)$ and such that : $i_2 \mid i_1$, $i_2 \mid (c - 2)$. Then there exists an ordinary Drinfeld A -module Φ over L of rank 2, such that: $L^\Phi \simeq M$.*

Lastly, we will make a statistic about the ordinary Drinfeld A -modules such that the A -modules L^Φ are cyclic, we note by $C(d, m, q)$ the proportion of the number (of isomorphisms of) ordinary Drinfeld A -modules, of rank 2 such that the A -modules structures L^Φ are cyclic, this means : if we note by $\#\{\Phi, \text{isomorphism, ordinary}\}$ the number of classes of L -isomorphisms of an ordinary Drinfeld Modules of rank 2, we have :

$$C(d, m, q) = \frac{\#\{\Phi, L^\Phi \text{ cyclic}\}}{\#\{\Phi, \text{isomorphism, ordinary}\}},$$

and we note by $C_0(d, m, q)$ the proportion of the number (of isogeny classes of) ordinary Drinfeld A -modules, of rank 2 such that the A -modules L^Φ are cyclic, otherwise, if we note by $\#\{\Phi, \text{isogeny, ordinary}\}$ the number of isogeny classes, of ordinary Drinfeld modules of rank 2, we have :

$$C_0(d, m, q) = \frac{\#\{\text{isogeny Classes of } \Phi, L^\Phi \text{ cyclic}\}}{\#\{\Phi, \text{isogeny, ordinary}\}}.$$

Of course, these numbers are depending on q and also on d, m . One of our important results is :

Proposition 1.3. $C(d, m, q) = C_0(d, m, q) = 1$ if and only if $m = d = 1$.

This means that, to have a Cyclic Drinfeld A -modules we must have a trivial extension L , we give also some values for $C(d, m, q)$ and $C_0(d, m, q)$ corresponding to some given values of d and m , for example :

Proposition 1.4. *We put $d = 2$ and $m = 1$. Let $H(O(D))$ the Hurwitz's number of classes for an order O which the imaginary determinant is D :*

$$C_0(2, 1, q) = \frac{q(q-1) - 5}{q(q-1) - 2},$$

$$C(2, 1, q) = \frac{q^3 - q^2 - q + 1 - [\frac{q-1}{2} \sum_{P_\Phi} \sum_{i_2, i_2^2 | 4-4\mu P} H(O(\frac{4-4\mu P}{i_2^2})) + (q-1) \sum_{P_\Phi} \sum_{i_2, i_2^2 | c^2-4\mu P} H(O(\frac{c^2-4\mu P}{i_2^2}))]}{q^3 - q^2 - q + 1}.$$

And we let think, in conjecture form, that for a big q the values of $C(d, m, q)$ and $C_0(d, m, q)$ will tend to 1.

2 Drinfeld Modules

Let E be an extension of $\mathbf{F}_q$, and let τ Frobenius of $\mathbf{F}_q$. We put $E\{\tau\}$ the polynomial ring in τ with the usual addition and the multiplication defined by:

$$\forall e \in E, \tau e = e^q \tau.$$

Definition 2.1. *Let R be the E -linearly polynomials set with the coefficient in E , that means that these elements are on the following form :*

$$Q(x) = \sum_{K \geq 0} l_K x^{q^K},$$

where $l_K \in E$ for every $K \geq 0$, and only a finite number of l_K is not null. The ring R is a ring by addition and the polynomial composition.

Lemma 2.1. $E\{\tau\}$ and R are isomorphic rings.

If we put $A = \mathbf{F}_q[T]$, $f(\tau) = \sum_{i=0}^v a_i \tau^i \in E\{\tau\}$ and $Df := a_0 = f'(\tau)$.
It is clear that the application :

$$E\{\tau\} \mapsto E$$

$$f \mapsto Df,$$

is a morphism of $\mathbf{F}_q$ -algebras.

Definition 2.2. An A -fields E is a field E equipped with a fix morphism $\gamma : A \longrightarrow E$. The prime ideal $P = \text{Ker}\gamma$ is called the characteristic of E . We say E has generic characteristic if and only if $P = (0)$; otherwise (i.e $P \neq (0)$) we said P is finite and E has finite characteristic.

We then have the following fundamental definition :

Definition 2.3. Let E an A -field and $\Phi : A \mapsto E\{\tau\}$ be homomorphism of algebra. Then Φ is a Drinfeld A -module over E if and only if :

1. $D \circ \Phi = \gamma$;
2. for some $a \in A$, $\Phi_a \neq \gamma(a)\tau^0$.

As was proved by Drinfeld in [6], such modules always exist.

Remark 2.1. 1. The above normalization is analogous to the normalization used in complex multiplication of elliptic curves. The last condition is obviously a non-triviality condition.

2. By Φ , every extension E_0 of E became an A -module by :

$$\begin{aligned} E_0 \times A &\rightarrow E_0, \\ (k, a) &\rightarrow k.a := \Phi_a(k). \end{aligned}$$

We will note this A -module by E_0^Φ .

Let $\overline{E}$ be a fix algebraic closure of E and Φ a Drinfeld module over E and I an ideal of A . As A is a Dedekind domain, one know that I may be generated by (at most) two elements $\{i_1, i_2\} \subset I$.

Since $E\{\tau\}$ has a right division algorithm, there exists a right greatest common divisor in $E\{\tau\}$. It is the monic generator of the left ideal of $E\{\tau\}$ generated by : Φ_{i_1} et Φ_{i_2} .

Definition 2.4. We set Φ_I to be the monic generator of the left ideal of $E\{\tau\}$ generated by Φ_{i_1} and Φ_{i_2} .

Definition 2.5. Let E_0 an extension of E and I an ideal of A . We define by : $\Phi[I](E)$ the finite subgroup of $\Phi[\overline{E_0}]$ given by the roots of Φ_I in $\overline{E}$.

If $a \in A$, then we set $\Phi[a] := \Phi[(a)]$. We can see it as :
 $\Phi[a] = \{ \text{set of roots of } \Phi(a) \text{ in } \overline{K} \}$, and $\Phi_I = \cap_{a \in I} \Phi[a]$. Then :

$$\Phi_a(\overline{E}) := \Phi[a](\overline{E}) = \{x \in \overline{E}, \Phi_a(x) = 0\}.$$

And for every ideal $Q \subset A$,

$$\Phi_Q(\overline{E}) := \Phi_Q(\overline{E}) = \cap_{a \in Q} \Phi_a(\overline{E}).$$

Remark 2.2. The groups : $\Phi[I](E)$ and $\Phi[I](\overline{E})$ are clearly stable under $\{\Phi_a\}_{a \in A}$.

Definition 2.6. Let Φ be a Drinfeld A -module over an A -field E . We say that Φ is supersingular, if and only if, the A -module constituted by a P -division points $\Phi_P(\overline{E})$ is trivial.

2.1 The Height and Rank of a Drinfeld Module Φ

Let Φ be a Drinfeld A -module over the A -field E . We note by $\deg_\tau$ the degree in indeterminate τ .

Definition 2.7. An element of $E\{\tau\}$ is called separable, if this constant coefficient is not null. It called purely inseparable if it is on the form $\lambda\tau^n$, $n > 1$ and $\lambda \in E$, $\lambda \neq 0$.

Let H be a global field of characteristic $p > 0$, and let ∞ one place (a Prime ideal) of H , we will note by H_∞ the completude of H at the place ∞ . We define the degree of function over A by :

Definition 2.8. Let $a \in A$, $\deg a = \dim_{F_q} \frac{A}{aA}$ if $a \neq 0$ and $\deg 0 = -\infty$.

We extend $\deg$ to K by putting $\deg x = \deg a - \deg b$ if $0 \neq x = \frac{a}{b} \in K$. If $A = \mathbf{F}_q[T]$, then the degree function is the usual polynomial degree. Let Q be a no null ideal of A , we define the ideal degrees of Q , noted $\deg Q$, by :

$$\deg Q = \dim_{F_q} \frac{A}{Q}.$$

Lemma 2.2. there exists a rational number r such that :

$$\deg_\tau \Phi_a = r \deg a.$$

Proof. It is easy to see that Φ is an injection, otherwise since $K\{\tau\}$ is an integre ring, $\text{Ker } \Phi$ is a prime ideal non null, so maximal in A and by consequence $\text{Im } \Phi$ is a field, so $\Phi = \gamma$. Since $-\deg_\tau$ define a no trivial valuation over $\text{Frac}(\Phi(A))$ (the fractions field of $\Phi(A)$) which is isomorphic to K , so $-\deg_\tau$ and $-\deg$ are equivalent valuations over K . There is rational number $r > 0$, such that :

$$r \deg = \deg_\tau.$$

□

Corollary 2.1. *Let $\Phi : A \mapsto E\{\tau\}$ be a Drinfeld A -module, so Φ is injective.*

Proposition 2.1. *The number r is a positive integer.*

Definition 2.9. *The number r is called the rank of the Drinfeld A -module Φ .*

For example if $A = \mathbf{F}_q[T]$, a Drinfeld A -module of rank r is on the form : $\Phi(T) = a_1 + a_2\tau + \dots + a_r\tau^r$, where $a_i \in E$, $1 \leq i \leq r-1$ and $a_r \in E^*$.

In this case $\text{char } E = P \neq (0)$ we can define the notion of height of a Drinfeld module Φ .

For this, we put $v_P : K \mapsto \mathbb{Z}$, an associate normalized valuation at P , this means, if $a \in K$ has a root over P of order t , we have $v_P(a) = t$.

For every $a \in A$, let $w(a)$ the most small integer $t > 0$, where τ^t occurs at Φ_a with a no null coefficient.

Lemma 2.3. *There exists a rational number h such that :*

$$w(a) = hv_P(a) \deg P.$$

Proposition 2.2. *The number h is a positive integer.*

Definition 2.10. *the integer h is called the height of Φ .*

For example if $A = \mathbf{F}_q[T]$, a Drinfeld A -module of height h of rank r is on the form : $\Phi(T) = a_0 + a_h\tau^h + \dots + a_r\tau^r$, where $a_i \in E$, $0 \leq i \leq r-1$ and $a_r \in E^*$.

Definition 2.11. *Let Φ and Ψ two Drinfeld A -modules over an A -field E and p an isogeny over E from Φ to Ψ .*

1. We say that p is separable if and only if $p(\tau)$ is separable.
2. We say that p is purely no separable if and only if $p(\tau) = \tau^j$ for one $j > 0$.

2.2 Norm of Isogeny

Definition 2.12. *Let F an integer over a ring A , with fractions field K . we note by $N_{K/K(F)}$ the determinant of the K -linearly application of multiplication by F to $K(F)$ (it is the usual norm if the extension $K(F)/K$ is separable.*

We can see that there is a morphism $N_{K/K(F)} : I_{\overline{A}} \rightarrow I_A$ from the fractional ideals groups of $\overline{A}$ to functionary ideals group of A , by this morphism we have:

Proposition 2.3. *The norm of isogeny is a principal ideal.*

Proposition 2.4. *Let $M_{fin}(A)$ the category of primes ideals of A and let $D(A)$ the monoïde of ideals of A . There exists an unique function :*

$$\chi : M_{fin}(A) \mapsto D(A),$$

multiplicative over the exact sequence and such that $\chi(0) = 1$ and $\chi(A/\wp) = \wp$ for every prime ideal $\wp$ of A .

Definition 2.13. *The function χ is called the Euler-Poincare characteristic.*

We can see $\chi(E^\Phi)$ and we note it by χ_Φ .

Proposition 2.5. *The ideals χ_Φ and P^m are principals (in A), and more clearly $\chi_\Phi = (P_\Phi(1))$ and $P^m = P_\Phi(0)$.*

1. We know that the norm of isogeny is a principal ideal, indeed $N(F) = P_\Phi(0)$ and $N(1 - F) = (P_\Phi(1))$ since F and $1 - F$ are a K -isogenys.
2. We can call χ_Φ the divisor of E -points, this divisor is analogue at the number of E -points for elliptic curves.
3. χ_Φ is the annihilator of A -module E^Φ . We can deduct that : $E^\Phi \subset (\frac{A}{\chi_\Phi})^r$.
4. The structure of A -module E^Φ is stable by the Frobenius endomorphism F .

Corollary 2.2. *If there are a Drinfeld A -module Φ , over a field E , of characteristic P and of degree m over A/P , then the ideal P^m is a principal ideal.*

Remark 2.3. *The above Corollary shows that there exists a restriction of the existence of Drinfeld A -modules.*

3 Drinfeld Modules over Finite Fields

We substitute the Field E , by L a finite extension of degree n of the finite field $\mathbf{F}_q$. Let $\tau : x \mapsto x^q$ be the Frobenius of $\mathbf{F}_q$, so the Frobenius F of L is $F = \tau^n$ and $\mathbf{F}_q[F]$ is the center of $L\{\tau\}$. We put $m = [L : A/P]$ and $d = \deg P$, then $n = m.d$. The function $-\deg$ define a valuation over K , the field of fractions of A . We put $r_1 = [K(F) : K]$ and r_2^2 the degree of the left field $\text{End}_L \Phi \otimes_A K$ over this center $K(F)$.

So a Drinfeld A -module Φ over L give a structure of A -module over the additive finite group L , this structure will be noted L^Φ . Let γ the application of A to L which an element a for A associate the constant of Φ_a , then it is easy to see that γ is a ring homomorphism, and that Φ and γ are equal over $A^* = \mathbf{F}_q^*$ the set of reversible elements of A .

Definition 3.1. *Let Φ be a Drinfeld A -module over a finite field L . We note by $M_\Phi(X)$ the unitary minimal polynomial of F over K .*

Proposition 3.1. *With the above notations : $M_\Phi(X)$ is an element of $A[X]$, equal to $P_\Phi^{\frac{1}{r_2^2}}$.*

Corollary 3.1. *For two Drinfeld A -modules Φ and Ψ , of rank r over a finite field L , then the following are equivalent :*

1. Φ and Ψ are isogenous,

2. $M_\Phi(X) = M_\Psi(X),$

3. $P_\Phi = P_\Psi.$

Proposition 3.2. *Let L be a finite extension of degree n over a finite field $\mathbf{F}_q$ and F the Frobenius of L . Then $L(\tau)$ is a central division algebra over $\mathbf{F}_q(F)$ of dimension n^2 .*

Definition 3.2. *Every $u \in L\{\tau\}$ can be writing on this form $u = \tau^h u'$ (since L is a perfect field) where $u' \in L\{\tau\}$ separable. The integer h is called the height of u and will be noted by $ht\ u$.*

In the finite field case, we can see the height of a Drinfeld A -module Φ over finite field L , the integer H_Φ , as been :

$$H_\Phi = \frac{1}{r} \inf\{ht\ \Phi_a, 0 \neq a \in P\}.$$

Remark 3.1. *It is easy to see that H_Φ is invariant under isogeny and that*

$$1 \leq H_\Phi \leq r.$$

Proposition 3.3. *Let Φ be a Drinfeld A -module of rank r over a finite field L , the following assertions are equivalent :*

1. *There exists a finite extension L' of L , such that the endomorphism ring $End_{L'}\Phi \otimes_A K$, has dimension r^2 over K .*
2. *Some power of the Frobenius F of L lies in A .*
3. *Φ is supersingular.*
4. *The field $K(F)$ has only one prime above P .*

Proposition 3.4. *Let Φ be a Drinfeld A -module of rank r and let Q be an ideal from A prime with P , then :*

$$\Phi_Q(\overline{L}) = \left(\frac{A}{Q}\right)^r.$$

Corollary 3.2. *Then we can deduct that : $\Phi_P(\overline{L}) = \left(\frac{A}{P}\right)^{r-H_\Phi}$.*

We can deduct from above mentioned Proposition the following important result, which characterize the supersingularity :

Proposition 3.5. *The Drinfeld A -module Φ is supersingular ($\Phi_P(\overline{L}) = 0$), if and only if, $r = H_\Phi$.*

Definition 3.3. *We say that the field L is so big if all endomorphism rings defined over $\overline{L}$ are also defined over L , i.e : $End_{\overline{L}}\Phi = End_L\Phi$.*

Two Drinfeld modules Φ and Ψ are isomorphic, if and only if, there exists an $a \in L$ such that : $a^{-1}\Phi_a = \Psi_a a$.

Lemma 3.1. *Let Φ be a Drinfeld A -module of rank r , over a finite field L , of characteristic P . The characteristic polynomial of Frobenius endomorphism F is :*

$$P_\Phi(X) = X^r + c_1 X^{r-1} + \dots + c_{r-1} X + \mu P^m, c_1, \dots, c_{r-1} \in A \text{ et } \mu \in \mathbf{F}_q^*.$$

Remark 3.2. *The fact that constant of the polynomial P is μP^m comes from the fact that $P_\Phi(0) = P^m$ in A .*

The following Proposition is an analogue of the Riemann's hypothesis for elliptic curves :

Proposition 3.6. *Let Φ be a Drinfeld A -module of rank r over finite field L which is a finite extension of degree n of $\mathbf{F}_q$. Then $\deg(w) = \frac{n}{r}$ for every root w of characteristic polynomial $P_\Phi(X)$.*

The following result is the Hasse-Weil's analogue for the elliptic curves :

Corollary 3.3. *Let $P_\Phi(X) = X^r + c_1 X^{r-1} + \dots + c_r X + \mu P^m$ be the characteristic polynomial of a Drinfeld Module Φ , of rank r , over a finite field L . Then:*

$$\forall 1 \leq i \leq r-1, \deg c_i \leq \frac{i}{r} m \deg P.$$

Proof. The proof can be deducted immediately by the above Proposition. $\square$

4 Drinfeld Modules of rank 2

In all next of this paper, Φ will be considered a Drinfeld A -module of rank 2, And $A = \mathbf{F}_q[T]$ for proof and more details see [1], [12] and [6].

4.1 Structure of A -module L^Φ

Let Φ be a Drinfeld A -module of rank 2, over a finite field L and let P this characteristic polynomial. About the A -module structure L^Φ , we have the following result :

Proposition 4.1. *The Drinfeld A -module Φ give a finite A -module structure L^Φ , which is on the form $\frac{A}{I_1} \oplus \frac{A}{I_2}$ where I_1 and I_2 are two ideals of A , such that: $\chi_\Phi = I_1 I_2$.*

Proof. Since the A -module L^Φ is a sub A -module of $\Phi(\chi_\Phi) \simeq \frac{A}{\chi_\Phi} \oplus \frac{A}{\chi_\Phi}$, then there are I_1 and I_2 in A such that : $L^\Phi \simeq \frac{A}{I_1} \oplus \frac{A}{I_2}$ and since the Euler-Poincare's Characteristic is multiplicative over the exact sequence we will have $\chi_\Phi = I_1 I_2$. $\square$

We put $I_1 = (i_1)$ and $I_2 = (i_2)$ (i_1 and i_2 two unitary polynomials in A).

Let $i = \text{pgcd}(i_1, i_2)$, it is clear by the Chinese lemma, that the non cyclicity of the A -module L^Φ , needs that I_1 and I_2 are not a prime between them, that means that $i \neq 1$, and since the relation $\chi_\Phi = I_1 I_2$, we will have : $i^2 \mid P_\Phi(1)$ ($\chi_\Phi = (P_\Phi(1))$).

In all the next of this paper, the condition above, will be considered verified, and more precisely we suppose that $I_2 \mid I_1$ (i.e : $i_2 \mid i_1$) otherwise L^Φ is a cyclic A -module and can be writing on this form A/χ_Φ .

Proposition 4.2. *If $L^\Phi \simeq \frac{A}{I_1} \oplus \frac{A}{I_2}$, then $i_2 \mid c - 2$.*

Proof. We know that the A -module structure L^Φ is stable by the endomorphisme Frobenius F of L . We choose a basis for A/χ_Φ , for which the A -module L^Φ will be generated by $(i_1, 0)$ and $(0, i_2)$.

Let $M_F \in \mathbf{M}_2(A/\chi_\Phi)$ the matrix of the endomorphisme Frobenius F in this basis. Then $M_F = \begin{pmatrix} a & b \\ a_1 & b_1 \end{pmatrix}$, where $a, b, a_1, b_1 \in A/\chi_\Phi$.

Although since : $\text{Tr } M_F = a + b_1 = c$ and $M_F(i_1, 0) = (i_1, 0)$ and $M_F(0, i_2) = (0, i_2)$, we will have $a.i_1 \simeq i_1 \pmod{\chi_\Phi}$ and then $a - 1$ is divisible by i_1 , of same for $b_1.i_2 \simeq i_2 \pmod{\chi_\Phi}$, that means that $b_1 - 1$ is divisible by i_2 and then: $c - 2 = a - 1 + b_1 - 1$ is divisible by i_2 (since we have always $i_2 \mid i_1$). $\square$

Let ρ be a prime ideal from A , different from the A -characteristic P , we define the finite A -module $\Phi(\rho)$ as been the A -module $(A/\rho)^2$.

The discriminant of the A -order: $A + g.O_{K(F)}$ is $\Delta.g^2$, where Δ is the discriminant of the characteristic polynomial $P_\Phi(X) = X^2 - cX + \mu P^m$. So each order is defined by this discriminant and will be noted by $O(\text{disc})$. It is clear, by the Propositions 4.1 that the inclusion $\Phi(\rho) \subset L^\Phi$ implies that $\rho^2 \mid P_\Phi(1)$ and $\rho \mid c - 2$. And if we note by $\text{End}_L \Phi$ the endomorphism ring of the Drinfeld A -module Φ , we have :

Proposition 4.3. *Let Φ be an ordinary Drinfeld A -module of rank 2, and let ρ an ideal from A different from the A -characteristic P of L , such that $\rho^2 \mid P_\Phi(1)$ and $\rho \mid c - 2$. Then $\Phi(\rho) \subset L^\Phi$, if and only if, the A -order $O(\Delta/\rho^2) \subset \text{End}_L \Phi$.*

To prove this Proposition we need the following lemma :

Lemma 4.1. *$\Phi(\rho) \subset L^\Phi$ is equivalent to $\frac{F-1}{\rho} \in \text{End}_L \Phi$.*

Proof. Since $L^\Phi = \text{Ker}(F - 1)$ and $\Phi(\rho) = \text{Ker}(\rho)$ (We confuse by commodity the ideal ρ with this generator in A) and we know by [3], Proposition 4.7.9, that for two isogenys, let by example $F - 1$ and ρ , we have $\text{Ker}(F - 1) \subset \text{Ker}(\rho)$, if and only if, there exists an element $g \in \text{End}_L \Phi$ such that $F - 1 = g.\rho$ and then $\Phi(\rho) \subset L^\Phi$, if and only if, $\frac{F-1}{\rho} = g \in \text{End}_L \Phi$. $\square$

We prove now the Proposition 4.3 :

Proof. Let $N(\frac{F-1}{\rho})$ the norm of the isogeny $\frac{F-1}{\rho}$, which is a principal ideal generated by $\frac{P_{\Phi}(1)}{\rho^2}$, and the trace (Tr) of this isogeny is $\frac{c-2}{\rho}$ then we can calculate the discriminant of the A -module $A[\frac{F-1}{\rho}]$ by:

$$\text{disc}A([\frac{F-1}{\rho}]) = \text{Tr}(\frac{F-1}{\rho})^2 - 4N(\frac{F-1}{\rho}) = \frac{c^2 - 4\mu P^m}{\rho^2} = \Delta/\rho^2 \Rightarrow$$

$$O(\Delta/\rho^2) \subset \text{End}_L \Phi.$$

We suppose now that : $O(\Delta/\rho^2) \subset \text{End}_L \Phi$ and we prove that $\Phi(\rho) \subset L^{\Phi}$. The Order corresponding of the discriminant Δ/ρ^2 is $A[\frac{F-1}{\rho}]$ this means that: $\frac{F-1}{\rho} \in \text{End}_L \Phi$ and so, by lemma 4.1 : $\Phi(\rho) \subset L^{\Phi}$. $\square$

Corollary 4.1. *If $O(\Delta/\rho^2) \subset \text{End}_L \Phi$, then L^{Φ} is not cyclic.*

Proof. We know that $\Phi(\rho)$ is not cyclic (since it is a A -module of rank 2), and then the necessary and sufficient conditions need for non cyclicity of A -module L^{Φ} are equivalent to the necessary and sufficient conditions to have $\Phi(\rho) \subset L^{\Phi}$. $\square$

We can so prove the following important Theorem :

Theorem 4.1. *Let $M = \frac{A}{I_1} \oplus \frac{A}{I_2}$, $I_1 = (i_1)$ And $I_2 = (i_2)$ such that : $i_2 \mid i_1$, $i_2 \mid (c-2)$. Then there exists an ordinary Drinfeld A -module Φ over L of rank 2, such that:*

$$L^{\Phi} \simeq M.$$

Proof. In fact, if we consider the Drinfeld A -module Φ , for which the characteristic of Euler-Poincare is giving by $\chi_{\Phi} = I_1 I_2$ and this endomorphism ring is $O(\Delta/i_2^2)$ where Δ is always the discriminant of the characteristic polynomial of the Frobenius F . We remind that $\Phi(\rho) \subset L^{\Phi}$ for every ρ an ideal A , different from P and verify $\rho^2 \mid P_{\Phi}(1)$ and $\rho \mid (c-2)$, if and only if, the A -order $O(\Delta/\rho^2) \subset \text{End}_L \Phi$. Let now $\rho = i_2$. Since by construction the A -order $O(\Delta/i_2^2) \subset \text{End}_L \Phi$ we have that $\Phi(i_2) \simeq (A/i_2)^2 \subset L^{\Phi}$. We know that L^{Φ} is included or equal to $\Phi(\chi_{\Phi}) \simeq \frac{A}{\chi_{\Phi}} \oplus \frac{A}{\chi_{\Phi}}$, we have so : $L^{\Phi} = \frac{A}{I_1} \oplus \frac{A}{I_2}$. $\square$

The Theorem above can be proved by using the following conjecture:

Conjecture 4.1. *Let $M \in \mathbf{M}_2(A/\chi_{\Phi})$, $\overline{P} = P \pmod{\chi_{\Phi}}$.*

We suppose : $(\det M) = \overline{P}^m$, $\text{Tr}(M) = c$ and $c \nmid P$. There exists a ordinary Drinfeld A -module over a finite field L of rank 2, for which the Frobenius matrix associated, is M_F , and such that :

$$M_F = M \in \mathbf{M}_2(A/\chi_{\Phi}).$$

We put the following matrix :

$$M_F = \begin{pmatrix} c-1 & i_1 \\ i_2 & -1 \end{pmatrix} \in \mathbf{M}_2(A/\chi_{\Phi}).$$

We can see that the three conditions of the conjecture are realized then there exists an ordinary Drinfeld A -modules Φ over L of rank 2, such that : $L^{\Phi} \simeq M$.

4.2 Deuring Theorem

The following Theorem, proved by Max-Deuring in [15] and [19] is used for the proof of the analogue of our principal result, in elliptic curves case :

Theorem 4.2. *Let E_0 be an elliptic curve over a finite field of characteristic p , with a no trivial endomorphism F_0 . Then there exists an elliptic curve E over a field of numbers and an endomorphisme F from E such E_0 is isomorphic to $\overline{E}$ and F_0 corresponding to F under this isomorphism.*

From the previous Theorem, we can deduct the following Theorem :

Theorem 4.3. *let $N \in \mathbb{N}$, $M = \begin{pmatrix} a & b \\ a_1 & b_1 \end{pmatrix} \in \mathbf{M}_2(\mathbb{Z}/N\mathbb{Z})$ and $\mathbf{F}_q$ a finite field with q elements, we suppose:*

1. $(\det M) = q \pmod{N}$;

2. $|a + b_1| \leq 2\sqrt{q}$.

There exists a Frobenius endomorphism F which verifies : $F^2 - cF + q = 0 \pmod{N}$, such that $c = a + b_1$ and this matrix $M_F \in \mathbf{M}_2(\mathbb{Z}/N\mathbb{Z})$ is exactly M .

This Theorem is used to prove the following Theorem :

Theorem 4.4. *Let $M = \begin{pmatrix} c-1 & -A \\ B & 1 \end{pmatrix} \in \mathbf{M}_2(\mathbb{Z}/N\mathbb{Z})$ and $\mathbf{F}_q$ is a finite field with q elements, such that : $|c| \leq 2\sqrt{q}$, $B \mid A$, $B \mid c-2$, $A.B = N = q+1-c$, we suppose : $(c, q) = 1$. Then there exists an ordinary elliptic curve E over $\mathbf{F}_q$, such that :*

$$E(\mathbf{F}_q) \simeq \mathbb{Z}/A \oplus \mathbb{Z}/B.$$

5 Cyclicity Statistics for the A -module L^Φ

In this section, we make a statistic about the Drinfeld Modules Φ of rank 2, whose the structures L^Φ are cyclic, for this, we define $C(d, m, q)$ as been the ration of the number of (isomorphism classes of) Drinfeld modules of rank 2 with cyclic structure L^Φ to the number of L -isomorphisms classes of ordinary Drinfeld modules of rank 2, noted by $\#\{\Phi, \text{isomorphism, ordinary}\}$:

$$C(d, m, q) = \frac{\#\{\Phi, L^\Phi \text{ cyclic}\}}{\#\{\Phi, \text{isomorphism, ordinary}\}},$$

As same, we define $N(d, m, q)$ as been the ration of the number of (isogeny classes of) Drinfeld modules of rank 2 with not cyclic structure L^Φ to the number of L -isomorphisms isogeny of ordinary Drinfeld modules of rank 2, noted by $\#\{\Phi, \text{isogeny, ordinary}\}$:

$$N(d, m, q) = \frac{\#\{\Phi, L^\Phi \text{ non cyclic}\}}{\#\{\Phi, \text{isogeny, ordinary}\}}.$$

We remark that : $0 \leq C(d, m, q), N(d, m, q) \leq 1$. Since the no cyclicity of the structure L^Φ needs the fact that $i^2 \mid P_\Phi(1)$ and $i_2 \mid (c-2)$, it is natural to introduce i (so i_2) in the calculus of $C(d, m, q)$ and $N(d, m, q)$.

We fix the characteristic polynomial P_Φ , this means that we fix the isogeny classes of Φ , and we define :

Definition 5.1. We note by $n(P_\Phi, i_2) = \#\{\Phi : L^\Phi = \frac{A}{(i_1)} \oplus \frac{A}{(i_2)}\}$.

Remark 5.1. The number $n(P_\Phi, i_2)$ is equal to the number of isomorphisms classes of Φ whole the A -module $L^\Phi \simeq \frac{A}{(i_1)} \oplus \frac{A}{(i_2)}$, in one isogeny classes, from where is coming the correspondence between Φ and i_2 .

For $n(P_\Phi, i_2)$ we have by the Theorem 3.1 :

Lemma 5.1. Let $P_\Phi(X) = X^2 - cX + \mu P^m$ be the characteristic polynomial of an ordinary Drinfeld A -module Φ of rank 2, and let i_2 be an unitary polynomial of A . Then if $i_2 \mid c-2$ we have : $n(P_\Phi, i_2) \geq 1$, else $n(P_\Phi, i_2) = 0$.

We can deduct :

Corollary 5.1. with the above notations :

$$\#\{\Phi, L^\Phi \text{ non cyclic}\} = \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid P_\Phi(1)} n(P_\Phi, i_2) \cdot \#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\},$$

$$\#\{\Phi, L^\Phi \text{ cyclic}\} = \sum_{P_\Phi} \sum_{i_2, i_2^2 \nmid P_\Phi(1)} n(P_\Phi, i_2) \cdot \#\{i_2, i_2^2 \nmid P_\Phi(1) \text{ and } i_2 \mid (c-2)\};$$

And if we note by $n_0(P_\Phi, i_2) = \#\{\text{isogeny classes of } \Phi : L^\Phi = \frac{A}{(i_1)} \oplus \frac{A}{(i_2)}\}$, we have :

$$n_0(P_\Phi, i_2) = 1.$$

We note now by $\#\{\Phi, \text{isogeny, ordinary}\}$ the number of isogeny classes, for an ordinary module Φ , then we define :

$$N_0(d, m, q) = \frac{\#\{\text{isogeny classes of } \Phi, L^\Phi \text{ not cyclic}\}}{\#\{\Phi, \text{isogeny, ordinary}\}},$$

the same for

$$C_0(d, m, q) = \frac{\#\{\text{isogeny Classes of } \Phi, L^\Phi \text{ cyclic}\}}{\#\{\Phi, \text{isogeny, ordinary}\}},$$

We can so announce the following lemma :

Lemma 5.2. *With the notations above, we have :*

$$\begin{aligned}
N_0(d, m, q) &= \frac{\#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\}}{\#\{\Phi, \text{isogeny, ordinary}\}}, \\
N(d, m, q) &= \frac{\sum_{P_\Phi} \sum_{i_2, i_2^2 \mid P_\Phi(1)} n(\Phi, i_2) \cdot \#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\}}{\#\{\Phi, \text{isomorphism, ordinary}\}}, \\
C_0(d, m, q) &= \frac{\#\{i_2, i_2^2 \nmid P_\Phi(1) \text{ et } i_2 \mid (c-2)\}}{\#\{\Phi, \text{isogeny, ordinary}\}}, \\
C(d, m, q) &= \frac{\sum_{P_\Phi} \sum_{i_2, i_2^2 \nmid P_\Phi(1)} n(\Phi, i_2) \cdot \#\{i_2, i_2^2 \nmid P_\Phi(1) \text{ and } i_2 \mid (c-2)\}}{\#\{\Phi, \text{isomorphism, ordinary}\}}, \\
\text{and } N(d, m, q) + C(d, m, q) &= 1, \quad N_0(d, m, q) + C_0(d, m, q) = 1.
\end{aligned}$$

The calculus of $\#\{\Phi, \text{isogeny, ordinary}\}$, for an ordinary A -module Φ , has been calculated in [3] and [4], as been :

Proposition 5.1. *Let $L = F_{q^n}$ and P the A -characteristic of L .*

We put $m = [L : A/P]$ and $d = \deg P$:

1. *m is odd and d is odd :*

$$\#\{\Phi, \text{isogeny, ordinary}\} = (q-1)(q^{\lfloor \frac{m}{2}d \rfloor + 1} - q^{\lfloor \frac{m-2}{2}d \rfloor + 1} + 1).$$

2. *m, d even:*

$$\#\{\Phi, \text{isogeny, ordinary}\} = (q-1)\left(\frac{(q-1)}{2}q^{\frac{m}{2}d} - q^{\frac{m-2}{2}d} + 1\right).$$

As for the number L -isomorphisms classes, we will need the following result, for the proof and more details see [9] :

Proposition 5.2. *Let L be a finite extension of degree n over $\mathbf{F}_q$, then the number of L -isomorphisms classes of a Drinfeld A -module of rank 2 over L is $(q-1)q^n$ if n is odd and $q^{n+1} - q^n + q^2 - q$ else.*

And to calculate the number of L -isomorphisms classes for an ordinary Drinfeld modules, we will need to calculate the number of L -isomorphisms classes for an supersingular Drinfeld modules and subtract it from the global number of L -isomorphisms classes, for this, we have by [10] :

Proposition 5.3. *Let L be a finite extension of n degrees over $\mathbf{F}_q$, then the number of L -isomorphisms classes of an supersingular Drinfeld A -module of rank 2, over L is $(q^{n_2} - 1)$, where $n_2 = \text{pgcd}(2, n)$.*

The calculus of $C(d, m, q)$ will be calculated in function of the values of d and m which are two major values to determinate c because $\deg c \leq \frac{m \cdot d}{2}$.

And to calculate the number of L -isomorphisms classes existing in each isogeny classes, we need the following Definition for more information, see [13]:

Definition 5.2. Let L be a finite extension of degree n over $\mathbf{F}_q$, we define $W(F)$ as been :

$$W(F) = \sum_{\Phi, F=\text{Frobenius}(\Phi)} \text{Weigh}(\Phi)$$

where :

$$\text{Weigh}(\Phi) = \frac{q-1}{\#\text{Aut}_L \Phi}.$$

$W(F)$ is the sum of weights(noted $\text{Weigh}(\Phi)$) of number of L -isomorphisms classes existing in each isogeny classes of the module Φ which the Frobenius is F .

And to calculate $\#\text{Aut}_L \Phi$ we have the following lemma :

Lemma 5.3. Let Φ be an ordinary Drinfeld A -module of rank 2, over a finite field $L = \mathbf{F}_{q^n}$, then : $\#\text{Aut}_L \Phi = q - 1$.

By the previous lemma, we can see that $\text{Weight}(\Phi) = \frac{q-1}{\#\text{Aut}_L \Phi} = 1$, that means :

Corollary 5.2. In the case of ordinary Drinfeld modules of rank 2, $W(F)$ is the number of L -isomorphisms classes existing in each isogeny classes.

Definition 5.3. Let D be an imaginary discriminant and let l a polynomial for which the square is a divisor of D and let $h(\frac{D}{l^2})$ the number of classes of the order for which the discriminant is $\frac{D}{l^2}$. We define the number of classes of Hurwitz for an imaginary discriminant D , noted $H(D)$ by :

$$H(D) = \sum_l \sum_{l^2|D} h\left(\frac{D}{l^2}\right).$$

For more definitions and information about the numbers classess of Hurwitz, see [13] and [20].

Lemma 5.4. If α is an integral element over A , for which $O = A[\alpha]$ is an A -order, then $\text{disc}(A[\alpha])$ is equal to the discriminant of the minimal polynomial of α .

What is interesting for us is the calculus of the $\text{disc}(A[F])$ and since $\text{disc}(A[F]) = \text{disc}(P_\Phi)$. To calculate the number of classes $W(F)$, we have the following result, for proof see [13].

Proposition 5.4. Let L be a finite extension of degree n of a field $\mathbf{F}_q$ and F the Frobenius of L , then:

$$W(F) = H(\text{disc}(A[F])).$$

It remains for us to calculate $n(\Phi, i_2)$:

Lemma 5.5. *Let P_Φ be the characteristic polynomial of an ordinary Drinfeld A -module of rank 2, over a finite field L such that $L^\Phi = \frac{A}{(i_1)} \oplus \frac{A}{(i_2)}$, and let Δ the discriminant of the characteristic polynomial of the Frobenius F , then :*

$$n(P_\Phi, i_2) = H(O(\Delta/i_2^2)).$$

Proof. We know that to have $L^\Phi = \frac{A}{(i_1)} \oplus \frac{A}{(i_2)}$, we have certainly :

$\Phi(i_2) \simeq (A/i_2)^2 \subset L^\Phi$, that is equivalent to say, by the Proposition 4.3, that the A -order $O(\Delta/i_2^2) \subset \text{End}_L \Phi$ where Δ is always the discriminant of the characteristic polynomial of F , P_Φ , and that :

$$n(P_\Phi, i_2) = H(O(\Delta/i_2^2)).$$

□

We can calculate some values of $C(d, m, q)$ for some d and m :

6 The case : $d = m = 1$

In this case $L = A/P = \mathbf{F}_q$, the A -module $L^\Phi = A/P$, so it is cyclic, that means that $C(1, 1, q) = 1$.

We can find this result by more explicit way :

Proposition 6.1. *Let $L = F_{q^n}$ and P the A -characteristic of L , $m = [L, A/P]$ and $d = \deg P$. We suppose $m = d = 1$. then :*

$$C(1, 1, q) = C_0(1, 1, q) = 1.$$

Proof. $P_\Phi(1) = 1 - c + \mu P^m = 1 - c + \mu P$, : $i_2^2 \mid P_\Phi(1) \Rightarrow i_2$ is a no null constant, then an element of $\mathbf{F}_q^*$, so $(i_2) = A$ and $\frac{A}{i_1} \oplus \frac{A}{i_2} = \frac{A}{i_1}$, then L^Φ is cyclic, that means that $C(1, 1, q) = 1 \Rightarrow N(1, 1, q) = 0$.

To calculate $\#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\}$, we must have i_2 an element of $\mathbf{F}_q^*$ and $\deg i_2 > 0 \Rightarrow$

$$\#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\} = 0 \text{ and then :}$$

$$N_0(1, 1, q) = \frac{\#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\}}{\#\{\Phi, \text{isogeny, ordinary}\}} = 0$$

$$\Rightarrow C_0(1, 1, q) = 1.$$

□

By more precise way, we can announce :

Theorem 6.1. *Let $L = F_{q^n}$ and P the A -characteristic of L , $m = [L, A/P]$ and $d = \deg P$. Then:*

$$C_0(d, m, q) = C(d, m, q) = 1 \Leftrightarrow m = d = 1.$$

Proof. We have just seen that $C_0(1, 1, q) = C(1, 1, q) = 1$.

Conversely and by the absurd : $m.d > 1$ (this means $m.d \geq 2$), we take for example $m = 1$ and $d = 2$.

To have $C_0(d, m, q) = C(d, m, q) = 1$, we must have

$\#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\} = 0$, what is not true, since if $c = aT + b$, where $a \in \mathbf{F}_q^*$ and $b \in \mathbf{F}_q$, it is sufficient to have an unitary i_2 and such that : $i_2 \mid (c-2)$, for this, we take : $i_2 = a^{-1}(c-2)$ this stay compatible with the fact that $a^{-2}(c-2)^2 \mid 1 - c + \mu P$, since there are many solutions for the equation in i_2 , i.e in a and b :

$$a^{-2}(c-2)^2 \mid 1 - c + \mu P \Rightarrow a^{-2}(aT + b - 2)^2 \mid 1 - aT - b + \mu(T^2 + pT + p_0),$$

which implies the equations : $2a^{-1}\mu(b-2) = \mu p_1 - a$ and $\mu[a^{-1}(b-2)]^2 = 1 - b + \mu p_0 \Rightarrow 2^{-1}[\mu p_1 - a] = 1 - b + \mu p_0$ this gives a values of a for each value of b , from where the many possibilities of i_2 , for example $i_2 = T - (\mu(2p_0 - p_1))^{-1}$, so it is more easy in the case which $m.d > 2$ to find an i_2 , such that : $i_2^2 \mid P_\Phi(1)$ and $i_2 \mid (c-2)$. $\square$

7 The case: $m = 1$ and $d = 2$

In this case $n = m.d = 2$, and $n_2 = 2 \Rightarrow \#\{\Phi, \text{ isomorphism, ordinary}\} = q^3 - q - (q^2 - 1) = q^3 - q^2 - q + 1$.

Proposition 7.1. *Let $L = F_{q^n}$ and P the A -characteristic of L , $m = [L, A/P]$ and $d = \deg P$. We suppose $m = 1$ and $d = 2$. Then:*

$$C_0(2, 1, q) = \frac{q(q-1) - 5}{q(q-1) - 2},$$

$$C(2, 1, q) = \frac{q^3 - q^2 - q + 1 - [\frac{q-1}{2} \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid 4-4\mu P} H(O(\frac{4-4\mu P}{i_2^2})) + (q-1) \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid c^2-4\mu P} H(O(\frac{c^2-4\mu P}{i_2^2}))]}{q^3 - q^2 - q + 1}.$$

Proof. We start by calculating :

$$\frac{\#\{i_2, i_2^2 \mid P_\Phi(1)\}}{\#\{\Phi; \text{isogeny}\}}.$$

For this, we distinguish between two cases, the case where $c = 2$ and the case where $c \neq 2$.

Then for $c = 2$: $i_2^2 \mid P_\Phi(1) \Rightarrow i_2^2 \mid \mu P^m - 1$ this implies that if we put $i_2 = T + j_2$, $j_2 \in \mathbf{F}_q$ and $P(T) = T^2 + p_1T + p_0$ where $p_1, p_0 \in \mathbf{F}_q$, are irreducibles, we will have $p_1 = 2j_2$ and $\mu p_0 - 1 = \mu j_2^2$ we will have then the equation $\mu[p_0 - \frac{p_1^2}{4}] = 1 \Rightarrow \mu(p_1^2 - 4p_0) = -4$, since the fact that $p_1^2 - 4p_0$ is not a square P is irreducible in A , we will have $-\mu$ no square, that means that

the number of possible μ is $\frac{q-1}{2}$, if we consider the fact that p_0, p_1 are fixe, we will have $\frac{(q-1)}{2}$ solutions, then it remains to calculate : for the case $c \neq 2$, for this, we put : $i_2 = T + j_2$, $j_2 \in \mathbf{F}_q$ and $c = aT + b$ where $a \in \mathbf{F}_q^*$ and $b \in \mathbf{F}_q$. The fact that $i_2 \mid (c-2) \Rightarrow j_2 = \frac{b-2}{a}$ and since $i_2^2 \mid P_\Phi(1)$ we will have : $1 - (aT + b) + \mu(T^2 + p_1T + p_0) = \mu(T + j_2)^2 \Rightarrow 1 - b + \mu p_0 = \mu j_2^2$ and $\mu p_1 - a = 2\mu j_2$, then : $\mu = \frac{a}{p_1 - 2j_2} = \frac{a}{p_1 - 2(\frac{b-2}{a})}$ and then :

$$\frac{a}{p_1 - 2(\frac{b-2}{a})}[p_0 - (\frac{b-2}{a})^2] + 1 - b = 0.$$

The solution numbers of this equation in (a, b) (p_0, p_1 are fixe) give us

$\#\{i_2, i_2^2 \mid P_\Phi(1) \text{ et } i_2 \mid (c-2)\}$, then we will have $(q-1)$ possible cases for i_2 . Then :

$$\begin{aligned} N_0(2, 1, q) &= \frac{\#\{i_2, i_2^2 \mid P_\Phi(1)\}}{\#\{\Phi, \text{ isogeny, ordinary}\}} \\ &= \frac{(q-1) + \frac{(q-1)}{2}}{(q-1)[(\frac{q-1}{2})q - 1]} \\ &= \frac{\frac{3(q-1)}{2}}{(q-1)[(\frac{q-1}{2})q - 1]} \\ &= \frac{3}{q(q-1) - 2} \Rightarrow \end{aligned}$$

$$\begin{aligned} C_0(2, 1, q) &= 1 - \frac{3}{q(q-1) - 2} \\ &= \frac{q(q-1) - 5}{q(q-1) - 2}. \end{aligned}$$

And for $N(2, 1, q)$:

$$\begin{aligned} N(2, 1, q) &= \frac{\sum_{P_\Phi} \sum_{i_2} n(\Phi, i_2) \cdot \#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\}}{\#\{\Phi, \text{ isomorphism, ordinary}\}} \\ &= \frac{\sum_{P_\Phi} \sum_{i_2, i_2^2 \mid 4-4\mu P} H(O(\frac{4-4\mu P}{i_2^2})) \cdot \frac{q-1}{2} + \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid c^2-4\mu P} H(O(\frac{c^2-4\mu P}{i_2^2}))(q-1)}{q^3 - q^2 - q + 1} \\ &= \frac{\frac{q-1}{2} \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid 4-4\mu P} H(O(\frac{4-4\mu P}{i_2^2})) + (q-1) \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid c^2-4\mu P} H(O(\frac{c^2-4\mu P}{i_2^2}))}{q^3 - q^2 - q + 1}. \end{aligned}$$

Finally : $C(2, 1, q) = 1 - N(2, 1, q) =$

$$1 - \frac{\frac{q-1}{2} \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid 4-4\mu P} H(O(\frac{4-4\mu P}{i_2^2})) + (q-1) \sum_{P_\Phi} \sum_{i_2, i_2^2 \mid c^2-4\mu P} H(O(\frac{c^2-4\mu P}{i_2^2}))}{q^3 - q^2 - q + 1} =$$

$$\frac{q^3 - q^2 - q + 1 - [\frac{q-1}{2} \sum_{P_\Phi} \sum_{i_2, i_2^2 | 4-4\mu P} H(O(\frac{4-4\mu P}{i_2^2})) + (q-1) \sum_{P_\Phi} \sum_{i_2, i_2^2 | c^2-4\mu P} H(O(\frac{c^2-4\mu P}{i_2^2}))]}{q^3 - q^2 - q + 1}. \quad \square$$

8 the case : $m = 2$ and $d = 1$

In this case $n = m.d = 2$, and $n_2 = 2 \Rightarrow \#\{\Phi, \text{isomorphism, ordinary}\} = q^3 - q - (q^2 - 1) = q^3 - q^2 - q + 1$.

Proposition 8.1. *Let $L = F_{q^n}$ and P the A -characteristic of L , $m = [L, A/P]$ and $d = \deg P$. We suppose that $m = 2$ and $d = 1$. Then:*

$$C_0(1, 2, q) = \frac{(q-1)q-4}{(q-1)q-2},$$

$$C(1, 2, q) = \frac{q^3 - q^2 - q + 1 - \sum_{P_\Phi} \sum_{i_2, i_2^2 | c^2-4\mu P} H(O(\frac{c^2-4\mu P}{i_2^2}))}{q^3 - q^2 - q + 1}.$$

Proof. We put $i_2 = T + j_2$, $j_2 \in \mathbf{F}_q$ and $P(T) = T + p$ where $p \in \mathbf{F}_q$. We start by calculate:

$$N_0(1, 2, q) = \frac{\#\{i_2, i_2^2 \mid P_\Phi(1)\}}{\#\{\Phi, \text{isogeny, ordinary}\}},$$

In this case $c = 2$, $i_2^2 \mid P_\Phi(1) = \mu P^2 - 1$ we will have : $2\mu p = 2\mu j_2 \mu p^2 - 1 = \mu j_2^2$ that means that $p = j_2$ and $\mu(p^2 - j_2^2) = 1$ so the contradiction, and then $\#\{i_2, i_2^2 \mid P_\Phi(1)\} = 0$. For $c \neq 2$ we calculate $\#\{i_2, i_2^2 \mid P_\Phi(1)$ and $i_2 \mid (c-2)\}$ and we remark first : $i_2 \mid (c-2) \Rightarrow j_2 = \frac{b-2}{a}$ and $i_2^2 \mid P_\Phi(1)$ implies that : $2p\mu - a = 2\mu j_2$ and $p^2 + 1 - b = \mu j_2^2$, finally we will have the equation :

$$p^2 + 1 - b = \frac{a}{2p - 2(\frac{b-1}{a})} (\frac{b-2}{a})^2$$

which is an equation in (a, b, p) and accept $(q-1)$ solutions, since the fact that p is fix, then :

$$\begin{aligned} N_0(1, 2, q) &= \frac{\#\{i_2, i_2^2 \mid P_\Phi(1)\}}{\#\{\Phi ; \text{isogeny, ordinary}\}} \\ &= \frac{q-1}{(q-1)((\frac{q-1}{2})q-1)} \\ &= \frac{1}{(\frac{q-1}{2})q-1}; \\ \Rightarrow C_0(1, 2, q) &= 1 - N_0(1, 2, q) \\ &= 1 - \frac{1}{(\frac{q-1}{2})q-1} \\ &= \frac{(q-1)q-4}{(q-1)q-2}. \end{aligned}$$

And for $N(1, 2, q)$:

$$\begin{aligned}
N(1, 2, q) &= \frac{\sum_{P_\Phi} \sum_{i_2} n(\Phi, i_2) \cdot \#\{i_2, i_2^2 \mid P_\Phi(1) \text{ and } i_2 \mid (c-2)\}}{\#\{\Phi, \text{ isomorphism, ordinary}\}} \\
&= \frac{\sum_{P_\Phi} \sum_{i_2, i_2^2 | c^2 - 4\mu P} H(O(\frac{c^2 - 4\mu P}{i_2^2}))(q-1)}{q^3 - q} \\
&= \frac{\sum_{P_\Phi} \sum_{i_2, i_2^2 | c^2 - 4\mu P} H(O(\frac{c^2 - 4\mu P}{i_2^2}))}{q(q+1)}.
\end{aligned}$$

In end:

$$\begin{aligned}
C(1, 2, q) &= 1 - \frac{\sum_{P_\Phi} \sum_{i_2, i_2^2 | c^2 - 4\mu P} H(O(\frac{c^2 - 4\mu P}{i_2^2}))}{q^3 - q^2 - q - 1} = \\
&= \frac{q^3 - q^2 - q + 1 - \sum_{P_\Phi} \sum_{i_2, i_2^2 | c^2 - 4\mu P} H(O(\frac{c^2 - 4\mu P}{i_2^2}))}{q^3 - q^2 - q + 1}.
\end{aligned}$$

□

8.1 $\lim_{q \rightarrow \infty} C_0(d, m, q)$ and $\lim_{q \rightarrow \infty} C(d, m, q)$ for $m.d \leq 2$

By the calculus of $C_0(d, m, q)$ and $C(d, m, q)$ for $m.d \leq 2$, we have :

Corollary 8.1. *Let $L = F_{q^n}$ and P the A -characteristic of L , $m = [L, A/P]$ and $d = \deg P$. Then :*

$$\lim_{q \rightarrow \infty} C_0(1, 1, q) = \lim_{q \rightarrow \infty} C_0(1, 2, q) = \lim_{q \rightarrow \infty} C_0(2, 1, q) = 1,$$

$$\lim_{q \rightarrow \infty} C(1, 1, q) = \lim_{q \rightarrow \infty} C(1, 2, q) = \lim_{q \rightarrow \infty} C(2, 1, q) = 1.$$

Proof. Since : $C_0(1, 2, q) = \frac{q(q-1)-4}{q(q-1)-2}$, $C_0(2, 1, q) = \frac{q(q-1)-5}{q(q-1)-2}$ et $C_0(1, 1, q) = 1$, where we can see that : $md \leq 2$, $\lim_{q \rightarrow \infty} C_0(d, m, q) = 1$. By other way, since $C_0(d, m, q) \leq C(d, m, q) \leq 1$ and by passing to the limit, we have : $\lim_{q \rightarrow \infty} C(d, m, q) = 1$, pour $md \leq 2$. □

By the results above, we can give the following conjecture :

Conjecture 8.1. *Let $L = F_{q^n}$ and P the A -characteristic of L , $m = [L, A/P]$ and $d = \deg P$. Then :*

$$\lim_{q \rightarrow \infty} C(d, m, q) = \lim_{q \rightarrow \infty} C_0(d, m, q) = 1.$$

8.2 Discussion and open Questions

By the last conjecture we can ask whether for a big values of d and m , we will have a cyclic modules ? otherwise :

$$\lim_{d \rightarrow \infty} C(d, m, q) = \lim_{d \rightarrow \infty} C_0(d, m, q) = 1?$$

$$\text{of same : } \lim_{m \rightarrow \infty} C(d, m, q) = \lim_{m \rightarrow \infty} C_0(d, m, q) = 1?$$

We can also ask whether the rank of a Drinfeld A -module Φ is decisive for the cyclicity of A -module L^Φ ?

Lastly, it is legitimate to ask if we can generalize the Theorem 4.1 for a Drinfeld A -modules Φ such that A is not $F_q[T]$ and such that the rank of Φ is more bigger than 2?

References

- [1] Bruno Angles. *Thèse de Doctorat : Modules de Drinfeld sur les corps finis*, Université Paul Sabatier-Toulouse III, no d'ordre 1872, (1994).
- [2] Bruno Angles. *One Some Subring of Ore Polynomilas Connected with Finite Drinfeld Modules*, J. Algebra 181 (1996) , no.2, 507–522.
- [3] M. S M. AHMED. Endomorphism Rings and Isogenies Classes for a Drinfeld A -Modules of Rank 2 over Finite Fields , preprint IML, 2004.
- [4] M. S M. AHMED. Modules de Drinfeld de rang 2 sur un corps fini, thèse de Doctorat, IML 2004.
- [5] David Goss. *Basic Structures of Function Field Arithmetic*, Volume 35 Ergbnise der Mathematik und ihrer Grenzgebiete, Springer.
- [6] V.G. Drinfeld. *Modules Elliptiques. Math*, USSR Sbornik, 94 (136), 594-627, 656, (1974).
- [7] V.G. Drinfeld. *Modules Elliptiques II Math*, USSR Sbornik, 102 (144), No 2, 182-194,325, (1977).
- [8] Ernst-Ulrich Gekeler. *On Finite Drinfeld Module*. J. Algebra 141, (1991), 187-203.
- [9] Ernst-Ulrich Gekeler and Brian A. Snyder. *Drinfled Modules Over Finite Fields* . Drinfeld Modules, Modular Schemes and Application. Aldenbiesen, (1996).
- [10] Igor Potemine. *Thèse de Doctorat : Arithmétiques des Corps Globaux de Fonctions et Géométrie des Schémas Modulaires de Drinfeld*, de l'Université Joseph Fourier(Grenoble I), (1997)
- [11] J.P. Serre. *Corps Locaux*, Hermann (1968).

- [12] Joseph. H. Silverman *The Arithmetic of Elliptic Curves*. Graduate Texts in Mathematics, 106.
- [13] J.K.Yu. *A Classe Number Relation Over Functions Fields*, J. Number Theory, 54, (1995), 318–340.
- [14] J-K. Yu. *Isogenis of Drinfeld Modules Over Finite Fields*, J. Number of Theory 54 (1995), no 1, 161–171.
- [15] M. Deuring. *Die Typen der Multiplikatorenringe Elliptischer Funktionenko-rper*, Abh. Math.sem.Univ.Hamburg, 14 (1941), 197-272.
- [16] M. A.Tsfasman-S. G. Vladut. *Algebraic-Geometric Codes*, Mathematics and Applications, Dordrecht et al, (1991).
- [17] R. Shoof. *Nonsingular Plane Cubic Curves Over Finite Filelds*, Journal of combinatory theory, series A 46, (1987), 183-211.
- [18] I. Reiner. *Maximal Orders*. Academic Presse, (1975).
- [19] H.G. Ruck. *A Note on Elliptic Curves Over Finite Fields*. Math. Comp. 49, no179, (1987), 301–304.
- [20] S.G. Vladut. *Cyclicity Statistics for Elliptic Curves Over Finite Fields*, Finite Fields Appl. 5 (1999), no 4, 354–363.
- [21] W. C. Waterhouse. *Abelian Varieties Over Finite Fields*. Ann. Sci. Ecole Norm. Sup2, (1969), 521-560.